SLAC-PUB-6625
18 August 1994
5 August 1997 (rev)

EGS4 in '97

A Decade Plus of Enhancements*

W. R. Nelson
Radiation Physics Department
Stanford Linear Accelerator Center
Stanford University, Stanford, California 94309, USA

A. F. Bielajew and D. W. O. Rogers
Institute for National Measurement Standards
National Research Council of Canada
Ottawa, K1A 0R6, Canada

H. Hirayama and Y. Namito
National Laboratory for High Energy Physics
1-1 Oho, Tsukuba-shi, Ibaraki-ken, 305, Japan

Presented at the First International Workshop on EGS4
26–29 August 1997, KEK, Tsukuba, Japan

[Originally presented at the World Congress on Medical Physics and Biomedical Engineering]
[21–26 August 1994, Rio de Janeiro, Brazil]

* Work supported by the Department of Energy, contract DE-AC03-76SF00515.

Organization of the Tutorial

- Introduction:
  - General remarks about Monte Carlo codes.
  - Quick history behind EGS.
- Description of the EGS4 Code System:
  - How it is organized and physics within it.
  - Basic features of the code.
  - Mechanics of running EGS4.
- Benchmarks. . . . . a necessity for credibility.
- *Additional features available after 1985*.

Recent Popularity of Monte Carlo Codes

Electron-photon codes have become very popular:

- Five-fold increase in journal papers (1983-88).
- Several good books on electron-photon Monte Carlo.
- MC codes often tend to be used as *black boxes*.

EGS4 has played a very direct role:

- Several M.S. and Ph.D. theses based on the code.
- Nine workshops have now been given on EGS4.
- Six Best Paper awards in *Medical Physics* journal.

Why the sudden interest?

Recent Monte Carlo Books on Radiation Transport

- I. Lux and L. Koblinger, Monte Carlo Particle Transport Methods: Neutron and Photon Calculations (CRC Press, 1991).
- R. L. Morin (Editor), Monte Carlo Simulation in the Radiological Sciences (CRC Press, 1988). [*Contributors:* H.-P. Chan, K. Doi, J. E. Goin, R. L. Morin, R. Nath, D. E. Raeside, J. C. Widman and J. F. Williamson]
- T. M. Jenkins, W. R. Nelson, A. Rindi, A. E. Nahum and D. W. O. Rogers (Editors), Monte Carlo Transport of Electrons and Photons (Plenum Press, 1988). [*Contributors:* P. Andreo, M. J. Berger, A. F. Bielajew, A. Del Guerra, B. Grosswendt, J. Halbleib, A. Ito, T. M. Jenkins, R. Mohan, A. E. Nahum, W. R. Nelson, D. W. O. Rogers, S. Seltzer and R. Wang]

Reasons for Increase in MC-code Popularity

- Analytic methods tend to be prohibitive.
- MC tends to be intuitive – appeals to experimentalists.
- Computers – faster and cheaper !!!

EGS4 Code in Particular

Powerful – Based on well-understood physics.

Versatile – General-purpose code.

Benchmarked – Extremely well-checked code.

Open Architecture – Many contributions by users.

User-friendly † – Reasonably well documented.

User-supported – Workshops, large user community.

Timely – A tool needed by medical physicists.

FREE !!! – Code readily available (*ftp*).

(† Maybe *"expert-friendly"* is a better description.)

Codes That Preceded EGS

Messel and Crawford code[1] – Australia (1958-1970).

- First to use computer for high-energy shower MC.
- Published excellent results – but code not available.

Zerby and Moran code[2] – ORNL (1962-1963).

- Motivated by the construction of SLAC.
- Excellent engineering calculations performed.
- Code not distributed outside ORNL.

Berger and Seltzer code[3] – NIST (1964-present).

- ETRAN – Excellent physics and MC techniques.
- User-friendly versions now available (ITS and MCNP).
- Unknown to the particle-physics community in 1966.

Nagel Code – University of Bonn (1963-1967)[4]

- Ph.D. dissertation (1964).
- Cylindrical geometry (only) — and hard coded!
- Only materials available were Pb and Cu.
- But. . . . . readily available (*e.g.*, DESY, MIT, SLAC).
- Brought to SLAC around 1966 (by Nagel himself).

DEVELOPMENT OF EGS3
(SLAC-HEPL Collaboration)

SHOWER code (by Nagel) became *seed code* for EGS3.

- Energy range extended (0.1 MeV to few GeV).
- Any of 100 elements (compounds, mixtures).
- PEGS3 code – easy way to make input data for EGS3.
- More efficient sampling than in Nagel's code.
- New processes were added.

Popularity of EGS3 in late 1970's linked to HE physics.

- For reasons given previously (versatile, credible, *etc.*).
- But also, perfect timing. . . . . *The November Revolution!*

Development of EGS4
(SLAC-KEK-NRCC Collaboration)

SLAC-KEK collaboration already underway in 1982:

- To fix bugs, extend flexibility for HE accelerators.

Rogers (NRCC) using EGS3 rather extensively:

- Tremendous low-energy benchmarking effort.
- Medical physics applications, detector responses.
- Importance of electron step-size revealed (ESTEPE).
- Bielajew and Rogers fix (low-energy) bugs.

SLAC-265 report issued December 1985[5].

PRESTA released in 1986 by Bielajew and Rogers[6]:

Major advance in electron transport algorithms!

Description of the EGS4 Code System

- EGS – *analog* Monte Carlo program
  - Actual physical processes simulated as closely as possible.
  - Variance reduction techniques not "built-in".
  - Good for studying *fluctuations* (*e.g.*, particle detectors).
  - Disadvantage: *Very time consuming.*
- Can introduce importance sampling via `WT` parameter:

```
CALL SHOWER(IQ,E,X,Y,Z,U,V,W,IR,WT)
```

  (normally `WT(NP)=1.0` by default in EGS4).
- PEGS code – created for efficiency reasons.

PEGS4 – Preprocessor for EGS4

- PEGS uses theoretical & empirical formulae.
  - Compute σ's, branching ratios, scattering coefficients. . . *etc.*
  - Output is a 'table' $\Longrightarrow$ for **very fast** look-up by EGS.
- Run PEGS code before running EGS
  - But only *once* for each medium.
  - Save PEGS output on disk for subsequent use by EGS.
- PEGS4 has other uses:
  - Diagnostic tool.
  - Calculate and plot cross sections, *etc.*
  - Check sampling routines via bootstrap technique.

Electron ($\pm$) Processes in EGS4

- Bremsstrahlung – $Z(Z+1)$.
  - $\theta_{\text{brem}} = mc^2/E$ (default).
  - Special θ_{brem}-sampling version available (macro).
- δ-ray production – Bhabha (e^+e^-) and Møller (e^-e^-).
- Collision loss – $dE/dx)_{\text{col}}$ (excitation/ionization)
  - Between discrete interactions.
  - Restricted $dE/dx)_{\text{col}}$ (*i.e.*, LET_Δ).
  - $dE/dx)_{\text{rad}}$ for soft x-rays (added to collision loss).
  - Density effect (Sternheimer-Berger-Seltzer) [revised by Hirayama in 1995[53]].
- Multiple scattering – Molière model.
- Positron annihilation – in-flight/at-rest.

Photon Processes in EGS4

- Pair production – $Z(Z+1)$.
  - $\theta_{\text{pair}} = mc^2/E$ (default).
  - Special θ_{pair}-sampling version available (macro).
- Compton scattering (unbound).
- Coherent (Rayleigh) scattering.
- Photoelectric effect (excitation energy deposited locally).
  - Special *fluorescence* version available.
  - PE-angle sampling (macro) also available.

Basic Capabilities of EGS4

- Dynamic energy range – several TeV down to:
  - 1 keV (photons).
  - 10 keV (electrons).
- Geometry routines (SUBROUTINE HOWFAR) available .
- Combinatorial Geometry (MORSE-CG) User Codes.
- Transport in $\vec{E}$ and $\vec{B}$ fields.

- <u>Weighting and biasing</u> techniques:
  - Variance reduction and importance sampling.
  - Splitting, path-length biasing, Russian roulette.
  - Leading-particle biasing.
- <u>Transport initiation</u> (`CALL SHOWER`) with:
  - Energy spectrum (*e.g.*, synchrotron radiation).
  - Spatial and/or angular distribution.
  - Pi-zero decay.

Mechanics of Running an EGS4 Job

- User must create **User Code** consisting of:
  - **MAIN** program — 'driver' code.
  - **HOWFAR** — geometry subprogram.
  - **AUSGAB** — scoring subprogram.
- In **MAIN** program:
  - `CALL HATCH` to read in media data created by PEGS4.
  - `CALL SHOWER` as many times as you want shower events.
- User Code follows specific set of rules:
  - Details (including an example) are in the User Manual.
  - Set of tutorial examples also provided.

Division between User-Interface and EGS4



EXPERIMENTAL BENCHMARKS*

- Many successful comparisons in high-energy physics.
- Most precise benchmarks come from medical physics.
- Accurate patient dosimetry must account for:
  - Scattering from machine components.
  - Scattering from inhomogeneities within human body (*i.e.*, bones, lungs,. . . the *interface effect* problem).

* For example, see Chapter 5 by Rogers and Bielajew in *The Dosimetry of Ionizing Radiation, Volume III*, K. R. Kase, B. E. Bjärngard and F. H. Attix (editors).

Schematic of a typical clinical accelerator used for radiotherapy



Heterogeneity benchmark experiment by Shortt *et al.* (1986)[46] using a monoenergetic point source of 20-MeV electrons. The dose per unit fluence was measured in a water tank containing both air and aluminum cylinders. Data was taken with a small solid-state detector and then normalized to a single point on the water-only curve. Experimental results (solid lines) are compared with EGS4 calculations (histograms).




Water only
(a)
Water with aluminum cylinder
(b)
Water with air cylinder
(c)
ABSORBED DOSE / FLUENCE (10-10 Gy-cm2)
DEPTH (cm)
2-91
6700A25


0.10 cm behind air cylinder
(a)
0.60 cm behind air cylinder
(b)
2.10 cm behind air cylinder
(c)
ABSORBED DOSE / FLUENCE (10-10 Gy-cm2)
RADIAL POSITION (cm)
2-91
6700A26

IMPROVEMENTS TO EGS4 SINCE 1985

- This will be a brief description only.
- Interested readers should consult references in

  A. F. Bielajew, H. Hirayama, W. R. Nelson and D. W. O. Rogers, "History, Overview and Recent Improvements in EGS4", NRC-PIRS-0436 (1994)

- Most improvements/enhancements are
  - supplied with UNIX-version distribution, *or*
  - obtained directly from author(s).
- For the most part, all changes are *options* to EGS4:
  - must be "switched on" via flags, *and/or*
  - by including "macros" in User Codes.

Improvements/enhancements fall into three groups

I. Changes to *Physics Modeling* in EGS4

PRESTA, Angle Sampling (Brem,Pair,PE), Fluorescence, EM-Fields, Polarization, Doppler Broadening, Compton Binding, Single Scattering, Cross-Section Improvement.

II. Development of *Tools* and *Techniques*

Forced Interactions, Range Rejection, Bremsstrahlung Splitting, Long-Sequence Random-Number Generation, PEGS Tools, Graphics Tools.

III. *Systems* and other *Support*

New Platforms (UNIX,PC), Listserv, Anonymous-ftp Timing Benchmark Database, Courses, User Groups.

PRESTA

PARAMETER REDUCED ELECTRON STEP TRANSPORT ALGORITHM[6]

PRESTA-I – Introduced three new aspects to transport of electrons (shortly after release of EGS4).

Path-Length Correction (IPLC) – A refined method for calculating average curvature between multiple scattering sub-steps (Standard-EGS4 overestimates by up to a factor of 2).

Lateral-Correction Algorithm (ILCA) – Introduces extra lateral component, correlating it to the multiple scattering angle at end of sub-step (Standard-EGS4 ignores this, underestimates lateral diffusion).

Boundary-Crossing Algorithm (IBCA) – Sub-steps made shorter in vicinity of boundaries (avoids transport artefacts near interfaces. . . . . but, Molière theory breaks down for small step sizes).

PRESTA (cont.)

PRESTA-II – A refinement of physics in PRESTA-I, making results much more stable with respect to step size.

- New multiple scattering theory – solves small step-size problem intrinsic to Molière theory.
- PLC and LCA much more step-size stable.
- An exact BCA – Boundaries are crossed in a purely analog mode (with respect to elastic scattering).
- Sub-step randomization – allows multiple boundaries to be crossed with minimal numerical artefact.

Bremsstrahlung and Pair Production Angular Distributions

- In the Standard-EGS4:
  - Bremsstrahlung and pair energies are sampled.
  - Polar angles are fixed at m/E and m/k, respectively.
- This thick-target approximation assumes that multiple Coulomb *scattering* "washes out" *production* angles.
- The rationalization for this can be shown by equating

$$\theta_{\text{Brem}} = \theta_{MS}$$
$$m/E \approx 15\sqrt{t}/E$$

resulting in $t \approx 0.001$ r.l.

- Originally only expected to be a problem for thin-targets.
- However, thick-target studies revealed:
  - Angular-distribution artefacts near central axis.
  - Occuring at both radiotherapy (MeV) and diagnostic (keV) energies.
- Bremsstrahlung-angle sampling macro/switch (`IBRDST`) was developed by Bielajew, Mohan and Chui[8] in 1989.
- Pair-angle sampling macro/switch (`IPRDST`) was also developed by Bielajew[9] in 1991 (motivated in this case by a high-energy physics experiment at SLAC).

PHOTOELECTRIC ANGLE SAMPLING

- In the Standard-EGS4:
  - Photoelectron given same direction as incident photon
- PE-angle sampling macro/switch (`IPHTER`) developed by Bielajew and Rogers (1986)[10].
  - Based on relativistic theory by Sauter.
  - Their comparision with low-energy TLD experiments showed no major effect.
  - Importance of this option recently demonstrated by *backscattered synchrotron radiation* studies for the new Asymmetric B-Factory project at SLAC.
- Sampling of *azimuthal* angle of photoelectron (from linearly-polarized PE interaction) by Namito *et al.* (1996)[47].

KAND L-SHELL FLUORESCENCE

- Standard-EGS4 does not create/transport fluorescent photons
- Substitute sampling routine `SUBROUTINE PHOTO` allows for generation of K_{α_1} and K_{β_1} fluorescent photons.
  - Originally developed by Nelson and Jenkins in 1985 – now standard (switch (`IEDGFL`) turns on fluorescence by region).
  - Requires auxiliary subroutine `EDGSET` (extended by K. Weaver (UCSF) to include all 100 elements). Hirayama *et al.* implemented 20 L-edge X-rays (1996)[48].
- Del Guerra *et al.* (1991)[11] have developed K and L-edge sampling scheme for compounds.
- Namito *et al.* implemented K-shell X-rays induced by electron-impact ionization (1996)[49].

Electromagnetic Field Transport

- `SUBROUTINE ELECTR` of Standard-EGS4:
  - Does *not* perform electron transport in $\vec{E}$ and $\vec{B}$ fields.
  - But... EGS4 contains *macros* and *templates* that allow the user to do this (call it hindsight!).
- A general theoretical treatment in

  A. F. Bielajew, "Electron Transport in $\vec{E}$ and $\vec{B}$ Fields", in Monte Carlo Transport of Electrons and Photons Below 50 MeV, (Plenum Press, 1988).

  describes how to accomplish this.

- Bielajew's EM-field treatise was preceded by a very innovative theoretical-experimental study:
  - Rawlinson, Bielajew, Munro and Galbraith (1984)[12].
  - Dose enhancement caused by *electric charge storage* in electron-irradiated phantoms.
  - Farrington Daniels award (best dosimetry paper in MP journal)!
- More recently Bielajew[13] has come up with another clever method for studying the benefits of employing strong longitudinal $\vec{B}$ fields to control the lateral spread of external electron (and photon) beams.

EGS_Windows_1/20 MeV Electrons in Water
Longitudinal B-Field (0 T) (slide #1 of 2)

EGS_Windows_1/20 MeV Electrons in Water
Longitudinal B-Field (20 T) (slide #2 of 2)

Cross Section Improvements

- PEGS4 Modifications – *Collision* Stopping Power
  - Duane, Bielajew and Rogers (1989)[14] added PEGS4 option (EPSTFL) for inputing arbitrary density-effect corrections.
  - PC program (EPSTAR) by Seltzer (1988)[15] was used—to calculate density-effect corrections for arbitrary materials (ICRU-37)[16].
  - Relatively small changes to the collision stopping power. . . but crucial for stopping-power-ratio studies.

- PEGS4 Modifications – *Radiative* Stopping Power
  - Rogers *et al.* (1989)[17] added PEGS4 option (`IAPRIM`) to make radiative stopping powers ICRU-37-compliant.
  - Used Seltzer's PC program called EPSA[15].
  - Option provides a global renormalization of $\frac{d\sigma}{dE})_{\text{brem}}$.
  - Noticeable changes in $\frac{d\sigma}{dE})_{\text{brem}}$ below 50 MeV—experimentally verified by Faddegon, Ross and Rogers (1990,1991)[18].
  - Namito *et al.* (1990)[19] observed significant changes at low energies—*e.g.*, 80-keV x-rays from Au—setting `IAPRIM=1` brought EGS4 into agreement with ETRAN.

CROSS SECTION IMPROVEMENTS (cont.)

- Photon cross sections (< 50 MeV) in Standard-EGS4:
  - Based on 1970 Storm-Israel data package (DLC-15)[20].
  - ANSI (ENDF/B-VI) recommends DLC-136/PHOTX for point-kernel and S_n-transport calculations.
  - Sakamoto (1993)[21] introduced PHOTX into PEGS4.
  - Essentially, different PE cross sections in PHOTX.
  - Small effects observed for exposure buildup factors.

Cross Section Change Along Path of Electron

- The *"fictitious-interaction"* sampling scheme (Standard-EGS4).
  - Next interaction sampled from σ_{tot} at beginning of step.
  - Charged particles lose energy continuously between discrete interactions and σ_{tot} is different at the end of its path.
  - If σ_{tot} decreases as E decreases, can use sampling trick.
  - Invalid assumption at low-E—as E decreases σ_{Moller} rises and overtakes the decrease in σ_{brem} (Rogers, 1984)[7].
  - Ma and Nahum (1992)[22] created a linear-variation model.
  - Recommend its use for $E < 1$ MeV and Møller creation thresholds below 20 keV (few % effects).

More Accurate Trigonometric Functions

- Tracking algorithms make frequent use of sines and cosines.
- To gain computational speed, Standard-EGS4 uses a look-up table macro in lieu of standard FORTRAN sine/cosine functions.
- If accurate small-angle modeling is crucial—it is easy to revert back to a sine-by-function macro (at the cost of CPU time).
- Li and Rogers (1993)[23], calculating electron mass scattering factors, found shortcomings with default look-up table macro.
- Li, Rogers, and Ma (1994)[24] created new table look-up macro with small-angle accuracy as well as speed (as high as 45%).

Single Elastic Scattering

- Standard-EGS4 uses Molière multiple-scattering theory.
  - Employs small-angle form of screened Rutherford cross section (couched in small-angle formalism).
  - Contains approximations that make angular distributions unstable for short electron sub-steps.
- Recently Bielajew, Wang and Duane (1993)[25]:
  - Modified EGS4 to allow for single elastic scattering (using partial-wave cross sections by Berger and Wang).
  - Purpose – to study Molière theory.
- Subsequent study by Bielajew (1994)[26] has resolved small step-size difficulty of Molière.

Binding Effect in the Compton Interaction

- Standard-EGS4 treats atomic electrons as "free" (electron binding important for $E_\gamma <$ few hundred keV).
- Electron binding manifests itself in three ways:
  - Reduction in total Compton-scattering cross section.
  - Modification of scattered photon angular distribution (*e.g.*, reduction in the forward direction).
  - Broadening of scattered photon energy spectrum.
- Including all three consistently is the best way to treat binding effect in Compton scattering.

- Hirayama and Trubey (1988)[27] – First to include electron binding in Compton cross section for EGS4.
  - Calculated buildup factors for 40–200 keV x-rays.
  - Bound Compton modeling shown to have noticeable effect, especially at lower energies.
- Namito and Hirayama (1991)[28] and Samili and Dupeursinge (1991)[29] – First to also modify Compton angular distribution in EGS4 for electron binding.
  - Used same method implemented in ETRAN-based codes (SANDYL, MCNP, and ACCEPT of the ITS series).

LINEAR-POLARIZED PHOTON SCATTERING

- Number of synchrotron radiation facilities growing rapidly.
- Increasing need to include polarized photon scattering in MC codes.
- Standard-EGS4 considers all particles to be unpolarized.
- Namito, Ban and Hirayama (1993)[30] – First to implement linearly-polarized photon scattering into EGS4.
  - For both Compton and Rayleigh processes.
  - Calculations compared with series of benchmark experiments performed at KEK Photon Factory.

Absorbed dose measurement of a mono-energetic ($k_0 = 30$ keV) photon beam using TLDs in a soft tissue equivalent phantom. Intensity monitored by free-air ion chamber. Linear polarization ($P = 0.84$) monitored by Be scattering foil and Ge detectors. Symbols are measurements and lines are EGS4 calculations (Namito, Ban and Hirayama (1993)[30]).





Compton Scattering with Doppler Broadening

- In addition to neglecting electron binding, Standard-EGS4 ignores motion of electrons in the atomic cloud.
- Compton-scattered photon spectrum is *broadened* by pre-collision motion of the electron.
- Namito, Ban and Hirayama (1994)[31] – First to include Doppler broadening with Compton scattering in EGS4.
- In fact...they simultaneously accounted for *electron binding*, *Doppler broadening*, and *linear polarization* in a complete and totally consistent way.
- Calculations were compared with another series of benchmark experiments performed at KEK Photon Factory.

Mono-energetic ($k_0 = 40$ keV) linearly-polarized ($P = 0.84$) photon beam scattered by sample (S). Scattered photons counted by Ge detectors located at $\theta = 90^\circ$ relative to beam. Intensity monitored by free-air ion chamber. Collimators (C) define beam and opening angle of detectors. EGS4 calculation (histograms) include full Compton-binding effects plus linear polarization. EGS4(S)=without Doppler, EGS4(CP)=with Doppler (Namito, Ban and Hirayama (1994)[31]).





Forced Photon Interactions

- As stated earlier, Standard-EGS4 is an analog MC code.
  - Particles may pass through geometry without interacting!
  - For photons passing through *thin targets*—very wasteful.
- Rogers and Bielajew (1984)[32] have successfully used the technique of *forcing photon interactions* to eliminate this waste in some EGS4 applications.
- Rogers *et al.* (1985, 1994)[33,34] have also refined this technique by creating non-interacting *"fictitious photons"*, a method that is sometimes called *weighting-in-lieu-of-absorption*.
- Forced-interaction biasing and weighting is well described in

  A. F. Bielajew and D. W. O. Rogers, "Variance-Reduction Techniques", in Monte Carlo Transport of Electrons and Photons Below 50 MeV, (Plenum Press, 1988).

Range Rejection

- Range rejection – another variance-reduction technique
  - Electrons that simply cannot reach a region of interest are discarded "on the spot".
  - Approximations involved (see previous reference book).
- Rogers *et al.* (1994)[33] have refined the technique by using very accurate restricted range tables—obtained by integrating restricted stopping powers supplied by PEGS4.
- Range rejection can be quite powerful—as much as a factor of four has been gained in ion-chamber response calculations by Bielajew, Rogers and Nahum (1985)[35].

Bremsstrahlung Splitting

- Splitting – another variance-reduction technique
  - Set N photons in motion to improve statistics.
  - Give each photon a weight of $1/N$.
- Bremsstrahlung splitting (`IBRSPL`) has been developed for EGS4 by Bielajew, Mohan and Chui (1989)[8] (as part of brem-angle sampling macros discussed earlier).
- Faddegon, Ross and Rogers (1990,1991)[18] employed this technique at radiotherapy energies using $N = 5-30$.
- Namito *et al.* (1990)[19] used N-values as high as 300 to study 80-keV x-ray production from Au targets.

PEGS-Related Tools

- Interactive tool recently developed by Karr and Bielajew (1993)[36] to further automate the use of PEGS.
  - Called (`PIF`) – Prepare Input File for PEGS.
  - Includes ICRU-37 stopping powers (described earlier).
  - Maintains database – 100 elements/over 300 compounds.
- EXAMIN – NRCC User Code to understand PEGS output.
- Limitation currently imposed on operation of PEGS4:
  - Can only create *one* set of data at a time.
  - "Workaround" available (N. Hammond, EDS-Scion/U.K.).

LONG-SEQUENCE RANDOM NUMBER GENERATORS

- Standard-EGS4 comes with two RNGs:
  - One specific to IBM mainframe architecture.
  - One based on same generator but coded for generic 32-bit 2's-complement integer arithmetic (*e.g.*, VAXs).
  - Sequence length (periodicity) of 2^{30} ($\sim 10^9$).
- Marsaglia *et al.* (1990)[37] long-period ($2^{144} \sim 10^{43}$) RNG now recommend for EGS4 (slightly slower than original RNG).
  - Machine independent/parallel-implementation adaptable.
  - Choose 10^9 independent sequences from initial conditions.

Graphics Tools

- Several general-purpose packages for graphics output of particle tracks and geometries of EGS4 simulations:
  - `SHOWGRAF` from SLAC by Cowan and Nelson (1987)[38].
  - `SHOW` from NRCC by Mangin and Bielajew.
  - `EGS_Windows_1` from NRCC by Wiebe and Bielajew (1991)[39].
  - `EGS_Windows_2` from LBL by Chatterjee and Donahue.
  - `EGS_Windows_3` from NRCC by Zurawski and Bielajew.
- Above packages have different functions and require specific hardware and software. See Bielajew (1993)[40] for details.
- Hirayama *et al.* have developed EGS4-graphics for the PC (1994)[50].

SHOWGRAF/SLAC 40-inch Bubble Chamber
Single 1-GeV Photon (20kG) (slide #1 of 4)

SHOWGRAF/SLAC 40-inch Bubble Chamber

Single 1-GeV Photon (20kG) – Closeup (slide #2 of 4)

EGS_Windows_1/Varian 2100C Accelerator (slide #3 of 4)

(Cover-page photo from SLAC Beam Line 23(1) (1993))

EGS_Windows_3/Varian 2100C Accelerator (slide #4 of 4)

Machine/System Distributions

- Originally, EGS4 was supported for only two machine types:
  - IBM/VM and VAX/VMS mainframes.
  - Example scripts (`EXEC` and `.COM` files) were provided.
  - Original distribution still available from SLAC or RSIC.
- Japanese-computer versions available from H. Hirayama (KEK).
- Around 1988 Walker *et al.* (1988,1990,1992)[41,42] volunteered to manage and distribute an IBM-PC version of EGS4.
- UNIX distributions:
  - Developed and maintained at NRCC by Bielajew (1990,1993)[43].
  - Includes most improvements/enhancements described here.
  - Macintosh distribution from KEK (1997)[51].

Timing Benchmark Database

- Best way to compare performance—*use one's own application*.
  - Standard timing benchmark code for radiotherapy created by Bielajew and Rogers (1992)[44].
  - For wide variety of computers—PCs to supercomputers.
  - Separate PC comparison (same code) by Walker *et al.* (1992)[42].
  - Latest combined results (by many contributors) maintained in *anonymous-ftp servers* (described below).
- Yasu *et al.* (1993)[45] compiled HE physics timing benchmarks.

LISTSERV AND ANONYMOUS-ftp SUPPORT

- *Listserv* – Electronic-mail discussion list (`EGS4-L`).
  - Promote discussion within growing EGS-User Community.
  - Users can post questions to this list that can be answered by the EGS-community-at-large.
  - To subscribe, send an e-mail message to:

    `listserv@slac.stanford.edu` (Internet)

    saying: `SUBscribe EGS4-L` "Your full name"
  - To post questions/comments/answers, send e-mail to:

    `egs4-l@slac.stanford.edu`

Anonymous ftp

- Distribution of UNIX version of EGS4 most conveniently done by *anonymous ftp*.
- Current sites: `nrcnet0.nrc.ca` [132.246.160.2]
  `academic.lbl.gov` [128.3.252.168]
- Anonymous ftp sites are dynamic — browse periodically.

  Contain EGS4 Codes System distribution, plus. . . . . .

  . . . graphics-support code.

  . . . contributions from users.

  . . . most recent timing benchmark studies.

  . . . PostScript reprints of EGS-related papers and reports (including many mentioned in this tutorial).

User Groups, Courses and the EGS Web Page

- User Groups
  - EGS4 User's Meeting in Japan

    Recently finished sixth (annual) conference at KEK. Approximately 70 participants were from outside of KEK.
  - EGS4-User Group of France

    Has approximately 25 members and produces an EGS4 newletter. E. Sartori (OECD/NEA Data Bank, Issy-les-Moulineaux) distributes a PC version of EGS4 thoughout France.

- Courses
  - We have given nine EGS4 courses—in Ottawa, Seattle, London, Capri and Montpellier.
  - Two workshops on BEAM, an EGS4 User Code specifically designed as a software package for simulating radiotherapy treatement units[52], have also been given.
- EGS Web Page
  - A copy of this paper (and many others) can be obtained directly from the EGS Web Page:

    `http://ehssun.lbl.gov/egs/egs.html`
  - The authors acknowledge the support of Rob Stewert (PNNL) and Rick Donahue (LBL), who graciously maintain the EGS Web Page on our behalf. Thanks!

REFERENCES

1. H. Messel and D. F. Crawford, Electron-Photon Shower Distribution Function, (Pergamon Press, 1970).
2. C. D. Zerby and H. S. Moran, "A Monte Carlo calculation of the three-dimensional development of high-energy electron-photon cascade showers", ORNL-TM-422 (1962); C. D. Zerby and H. S. Moran, "Studies of the longitudinal development of high-energy electron-photon cascade showers in copper", ORNL-3329 (1962); C. D. Zerby and H. S. Moran, "Studies of the longitudinal development of electron-photon cascade showers", J. Appl. Phys. **34** (1963) 2445.
3. S. M. Seltzer, "An Overview of ETRAN Monte Carlo Methods", in Monte Carlo Transport of Electrons and Photons Below 50 MeV, T. M. Jenkins, W. R. Nelson, A. Rindi, A. E. Nahum and D. W. O. Rogers (editors) (Plenum Press, 1988).
4. H. H. Nagel and C. Schlier, "Berechnung von Elektron-Photon-Kaskaden in Blei für eine Primärenergie von 200 MeV", Z. Physik **174** (1963) 464; H. H. Nagel,"Elektron-Photon-Kaskaden in Blei: Monte-Carlo-Rechnungen für Primärelektronenergien zwischen 100 und 1000 MeV", Z. Physik **186** (1965) 319.
5. W. R. Nelson, H. Hirayama and D. W. O. Rogers, "The EGS4 Code System", SLAC-265 (1985).
6. A. F. Bielajew and D. W. O. Rogers, "PRESTA: The Parameter Reduced Electron-Step Transport Algorithm for Electron Monte Carlo Transport", NRC-PIRS-0042 (1986) and Nucl. Instr. Meth. **B18** (1987) 165.
7. D. W. O. Rogers, "Low energy electron transport with EGS", Nucl. Inst. Meth. **227** (1984) 535.
8. A. F. Bielajew, R. Mohan and C. S. Chui, "Improved bremsstrahlung photon angular sampling in the EGS4 code system", NRC-PIRS-0203 (1989).
9. A. F. Bielajew, "Improved angular sampling for pair production in the EGS4 code system", NRC-PIRS-0287 (1991).
10. A. F. Bielajew and D. W. O. Rogers, "Photoelectron angular distribution in the EGS4 code system", NRC-PIRS-0058 (1986).

11. A. Del Guerra, W. R. Nelson and P. Russo, "A simple method to introduce K-edge sampling for compounds in the code EGS4 for X-ray element analysis", Nucl. Instr. Meth. **A306** (1991) 378.

12. J. A. Rawlinson, A. F. Bielajew, P. Munro and D. M. Galbraith, "Theoretical and experimental investigation of dose enhancement due to charge storage in electron-irradiated phantoms", Med. Phys. **11** (1984) 814.

13. A. F. Bielajew, "The effect of strong longitudinal magnetic fields on dose deposition from electron and photon beams", Med. Phys. **20** (1993) 1171.

14. S. Duane, A. F. Bielajew and D. W. O. Rogers, "Use of ICRU-37/NBS Collision Stopping Powers in the EGS4 System", NRC-PIRS-0177 (1989).

15. S. M. Seltzer, "A PC-based program EPSTAR/ESPA," private communication (1988).

16. ICRU,"Stopping Powers for Electrons and Positrons", Report 37 (1984).

17. D. W. O. Rogers, S. Duane, A. F. Bielajew and W. R. Nelson, "Use of ICRU-37/NBS radiative stopping powers in the EGS4 system", NRC-PIRS-0177 (1989).

18. B. A. Faddegon, C. K. Ross and D. W. O. Rogers,"Forward Directed Bremsstrahlung of 10–30 MeV Electrons Incident on Thick Targets of Al and Pb", Med. Phys. **17** (1990) 773; B. A. Faddegon, C. K. Ross and D. W. O. Rogers, "Angular distribution of bremsstrahlung from 15 MeV electrons incident on thick targets of Be, Al and Pb", Med. Phys. **18** (1991) 727.

19. Y. Namito, W. R. Nelson, S. M. Seltzer, A. F. Bielajew and D. W. O. Rogers, "Low-energy x-ray production studies using the EGS4 code system", Med. Phys. **17** (1990) 557 (abstract).

20. E. Storm and H. I. Israel, "Photon Cross Sections from 1 keV to 100 MeV for Elements Z=1 to Z=100", Atomic Data and Nuclear Data Tables (1970) 565.

21. Y. Sakamoto, "Photon Cross Section Data PHOTX for PEGS4 Code", in "Proceedings of the Third EGS4 User's Meeting in Japan" (KEK, Japan) (1993) 77.

22. C.-M. Ma and A. E. Nahum, "A new algorithm for EGS4 low-energy electron transport to account for the change in discrete interaction cross-section with energy", Nucl. Instr. Meth. **B72** (1992) 319.

23. X. A. Li and D. W. O. Rogers, "Electron Mass Scattering Powers Calculated Using Monte Carlo Simulation", submitted to Med. Phys. (May, 1993).

24. X. A. Li, D. W. O. Rogers and C. M. Ma, "Rapid calculation of sines and cosines: an improved algorithm for EGS4", to be submitted to Nucl. Instr. Meth. (1994).

25. A. F. Bielajew, R. Wang and S. Duane, "Incorporation of single scattering in the EGS4 Monte Carlo code system: Tests of Moliere theory", Nucl. Inst. Meth. **B82** (1993) 503.

26. A. F. Bielajew, "Plural and multiple small-angle scattering from a screened Rutherford cross section", Nucl. Instr. Meth. **B86** (1994) 257.

27. H. Hirayama and D. K. Trubey, "Effects of incoherent and coherent scattering on the exposure buildup factors of low-energy gamma rays", Nucl. Sci. Eng. **99** (1988) 145.

28. Y. Namito and H. Hirayama, "Improvement of low energy photon transport calculation by EGS4 – electron bound effect in Compton scattering," Japan Atomic Energy Society, Osaka 401 (1991).

29. A. Samili and C. Depeursinge, "Adaptation du code EGS: Implémentation de l'effet de liaison des électrons lors d'une diffusion Compton", Swiss Federal Institute of Technology, March (1991).

30. Y. Namito, S. Ban and H. Hirayama, "Implementation of linearly-polarized photon scattering into the EGS4 code", Nucl. Instr. Meth. **A322** (1993) 277.

31. Y. Namito, H. Hirayama and S. Ban, "Implementation of Doppler broadening of Compton-scattered photons into the EGS4 code", Nucl. Instr. Meth. **A349** (1994) 489.

32. D. W. O. Rogers and A. F. Bielajew, "The Use of EGS for Monte Carlo Calculations in Medical Physics", NRC-PXNR-2692 (1984).

33. D. W. O. Rogers, B. A. Faddegon, G. X. Ding, C.-M. Ma, J. Wei and T. R. Mackie, "BEAM: A Monte Carlo code to simulate radiotherapy treatment units", submitted to Medical Physics (1994).

34. D. W. O. Rogers, G. M. Ewart, A. F. Bielajew and G. van Dyk, "Calculation of Contamination of the ^{60}Co Beam from an AECL Therapy Source", NRC-PXNR-2710 (1985).

35. A. F. Bielajew, D. W. O. Rogers and A. E. Nahum, "Monte Carlo simulation of ion chamber response to ^{60}Co – Resolution of anomalies associated with interfaces", Phys. Med. Biol. **30** (1985) 419.

36. J. L. Karr and A. F. Bielajew, "46PIF [Prepare Input File (for 46PEGS4)]", NRC-PIRS-0365 (1993).

37. G. Marsaglia, A. Zaman and W. W. Tsang, "Toward a Universal Random Number Generator", Stat. Prob. Lett. **8** (1990) 35.

38. R. F. Cowan and W. R. Nelson, "Producing EGS4 Shower Displays with the Unified Graphics System", SLAC-TN-87-3 (1987).

39. A. F. Bielajew and P .E. Wiebe, "EGS-Windows – A Graphical Interface to EGS", NRC-PIRS-0274 (1991).

40. A. F. Bielajew, "Graphics!," NRC-PIRS-0397 (1993).

41. S. Walker and D. Jette, "Installation of the EGS4 Monte Carlo code on a microcomputer system", Phys. Med. Biol. **33** Supplement 1 (abstract) (1988) 137; S. Walker and D. Jette, "Full installation of EGS4 Monte Carlo code on an 80386 microcomputer completed", Med. Phys. **17** (abstract) (1990) 511.

42. S. Walker, A. F. Bielajew, M. Hale and D. Jette, "Installation of EGS4 Monte Carlo code on an 80386-based microcomputer", Med. Phys. **19** (1992) 305.

43. A. F. Bielajew, "The EGS4 Monte Carlo code system on a SUN/UNIX platform", Med. Phys. **17** (abstract) (1990) 522; A. F. Bielajew, "How to manage the EGS4 system", NRC-PIRS-0391 (1993); A. F. Bielajew, "Running EGS4 on other machines", NRC-PIRS-0392 (1993).

44. A. F. Bielajew and D. W. O. Rogers, "A standard timing benchmark for EGS4 Monte Carlo Calculations", Med. Phys. **19** (1992) 303.

45. Y. Yasu, S. Ichii, S. Yashiro, H. Hirayama, A. Kokufuda and E. Suzuki, "High Energy Physics (HEP) Benchmark Program", KEK Preprint 93-155 (1993).

46. K. R. Shortt, C. K. Ross, A. F. Bielajew and D. W. O. Rogers, "Electron beam dose distributions near standard inhomogeneities", Phys. Med. Biol. **31** (1986) 235.

47. Y. Namito, S. Ban and H. Hirayama, "Photoelectron polarimetry simulation using EGS4 code", Japan Atomic Energy Society Fall Meeting, Tohoku University (1996).
48. H. Hirayama, Y. Namito and S. Ban, "Implementation of an L-shell photoelectron and an L X-ray for elements into the EGS4 code", in "Proceedings of the Sixth EGS4 User's Meeting in Japan" (KEK, Japan), KEK Proceedings 96-10 (1996).
49. Y. Namito and H. Hirayama, "Implementation of the electron impact ionization into the EGS4 code", "Proceedings of the Sixth EGS4 User's Meeting in Japan" (KEK, Japan), KEK Proceedings 96-10 (1996).
50. H. Hirayama, Y. Namito, S. Ban, R. Ikeda and Y. Tokuda, "EGS4 shower display system, EGS4PICT(2), Windows version", KEK Internal 94-10 (1994); also see "EGS4PICT, Windows version 2.0" for quasi-interactive version, KEK Internal 96-9 (1996).
51. Y. Namito, H. Nakamura and H. Hirayama, Lecture Note, Installation of EGS4, KEK Internal 97-7 (1997).
52. D. W. O. Rogers, B. A. Faddegon, G. X. Ding, C.-M. Ma, J. Wei and T. R. Mackie, "BEAM: A Monte Carlo code to simulate radiotherapy units", Med. Phys. **22** (1995) 503.
53. H. Hirayama, "Revision of the Sternheimer density effect coefficients in PEGS4", KEK Internal 95-17 (1995).
54. The best public discussion of the new physics in PRESTA-II can be found on the World Wide Web at: `http://gold.sao.nrc.ca/inms/irs/people/abielajew/pubs/proceedings/ANS-MnC_97.ps`