TRANSLATED FOR
STANFORD LINEAR ACCELERATOR CENTER

NONLOCAL QUANTUM FIELD THEORY, NONLINEAR INTERACTION LAGRANGIANS,
AND CONVERGENCE OF THE PERTURBATION-THEORY SERIES

by

G.V. Efimov

It is shown within the nonlocal quantum theory of one-component scalar field ϕ , that for an essentially nonlinear interaction Lagrangians $L_I(\phi)$, such that $L_I(\phi) \rightarrow 0$ at $\phi \rightarrow \pm \infty$, the nonlocal form factor can be chosen so that S -matrix will be finite, unitary and macrocausal in each order of perturbation theory, and the whole perturbation series will converge absolutely in the Euclidean region.

Only a few years ago, essentially nonlinear interaction Lagrangians for quantized fields were regarded as an exotic phenomenon in field theory, and their study was of purely academic interest. Recently, however, the requirement of invariance with respect to the chiral group, and study of nonlinear representations of this group, have led to a direct physical interpretation of nonlinear chiral-invariant Lagrangians (see, for example, ref. 1). The problem then arose as to how to operate with these Lagrangians. The initial idea was essentially connected with the desire to depart from the quantum field theory difficulties in the

analysis of this type of Lagrangian by considering them essentially as phenomenological and carrying out calculations only in the first nonvanishing orders in the interaction Lagrangian, completely neglecting contributions from the so-called closed loops².

This seems to us to be inconsistent; it can be regarded only as a first step in the study of nonlinear Lagrangians. On the other hand, there have already been a number of attempts to formulate a perturbation theory with an essentially nonlinear interaction^{3,4}. Although these cannot yet be regarded as final formulations either, they are, nevertheless, attempts to develop a realistic instrument for investigating the S-matrix in higher-orders of perturbation theory.

The first papers on the application of methods involving essentially nonlinear Lagrangians in the theory of chiral symmetries have now begun to appear⁵.

We shall show in this paper that within the framework of nonlocal quantum field theory with essentially nonlinear interaction Lagrangians, the S-matrix is not only finite and unitary in each order of perturbation theory, but also the entire series converges absolutely for a certain definite class

of interaction Lagrangians.

1. Formulation of the problem

It has been shown⁶ that the nonlocal unitary theory of a one-component scalar field can be constructed as follows:

Physical hypothesis: Nothing can be said in principle about the behavior of particles at sufficiently small space-time distances.

Mathematically, this means that the propagator for the free scalar particle

$$\Delta_0(x_1 - x_2) = \langle 0 | T(\phi(x_1) \phi(x_2)) | 0 \rangle \quad (1.1)$$

can be augmented by a generalized function from a suitable class of nonlocal generalized functions⁷ because Eq. (1.1) is undefined in the small space-time region around the point $x_1 = x_2$. Instead of the propagator $\Delta_0(x_1 - x_2)$ we have

$$D_0(x_1 - x_2) = \Delta_0(x_1 - x_2) + \frac{1}{i} K(x_1 - x_2), \quad (1.2)$$

where

$$K(x_1 - x_2) = \sum_{n=0}^{\infty} \frac{c_n}{(2n)!} \square^n \delta^{(4)}(x_1 - x_2). \quad (1.3)$$

The quantity $K(x_1 - x_2)$ is a generalized function from a suitable class of nonlocal generalized functions⁷. In momentum-space we obtain:

$$i \tilde{D}_0(p^2) \approx \frac{1}{m^2 - p^2 - i\epsilon} + \tilde{K}(\ell^2 p^2) \approx \frac{\tilde{V}(-\ell^2 p^2)}{m^2 - p^2 - i\epsilon}, \quad (1.4)$$

where ℓ can be interpreted as an elementary length.

It is shown in refs. 6 and 7 that if:

$$\text{I} \quad \overline{\lim}_{|z| \rightarrow \infty} \frac{\ell_n \ell_n |\tilde{V}(|z|)|}{\ell_n |z|} < 1. \quad (1.5)$$

$$\text{II} \quad \tilde{V}(-z) = o\left(\frac{1}{z}\right) \quad \text{at } z \rightarrow -\infty. \quad (1.6)$$

$$\text{and III} \quad \tilde{V}(-\ell^2 m^2) = 1, \quad (1.7)$$

then there exists a singular procedure for a limiting transition such that it is possible to transform to the Euclidean metric of space-like external momenta or space-like co-ordinates. The S-matrix in the Euclidean metric turns out to be finite in each order of the perturbation theory. Transition to the physical region is achieved by analytic continuation with respect to the invariant momentum variables. It turns out that the S-matrix is unitary in each order of the perturbation theory^{6,7}.

In the Euclidean formulation the S-matrix can be represented by the following formal expression

$$S[\phi] = \exp \left\{ \frac{1}{2} \iint d^4 x_1 d^4 x_2 D_0(x_1 - x_2) \frac{\delta^2}{\delta \phi(x_1) \delta \phi(x_2)} \right\} \times \\ \times \exp \left\{ g \int d^4 x U(\phi(x)) \right\}, \quad (1.8)$$

$$D_0(x) = \frac{1}{(2\pi)^4} \int \frac{d^4 k \tilde{V}(\ell^2 k^2)}{m^2 + k^2} e^{ikx}. \quad (1.9)$$

In this representation the S-matrix is regarded as a functional of the real field ϕ (not an operator). The entire integration in Eqs. (1.8) and (1.9) is carried out over the four-dimensional Euclidean space x and k .

If we expand in terms of the interaction constant g , we obtain

$$S[\phi] = \sum_{n=0}^{\infty} \frac{g^n}{n!} \int d^4 x_1 \dots \int d^4 x_n S_n(x_1, \dots, x_n), \quad (1.10)$$

$$S_n(x_1, \dots, x_n) = \exp \left\{ \frac{1}{2} \sum_{i,j=1}^n D_0(x_i - x_j) \frac{\delta^2}{\delta \phi(x_i) \delta \phi(x_j)} \right\} \prod_{j=1}^n U(\phi(x_j)), \quad (1.11)$$

In field theory the interaction Lagrangian

$$L_I(x) = g U(\phi(x)) \quad (1.12)$$

is usually taken in the normal form. This means that the diagonal terms are omitted in the sum of the differential operators in the index of the exponential in Eq. (1.11), i.e. it is assumed that

$$D_0(0) = 0. \quad (1.13)$$

The immediate reason for this choice is that in the usual local theory the quantity $D_0(0)$ is defined by a divergent integral and has therefore no meaning.

If the interaction Lagrangian given by Eq. (1.12) is a polynomial of the form $U(\phi) = \phi^m$, the operator in Eq. (1.11) is well defined in each order in g . However, nothing can be said about the properties of this series.

In this paper we shall investigate the existence of the functions $S_n(x_1, \dots, x_n)$ in Eq. (1.11) and of the entire functional $S[\phi]$ as a whole, depending on the form of the interaction Lagrangian given by Eq. (1.12).

We shall introduce two further important assumptions. First, we shall assume that the form factor $\tilde{V}(z)$ in Eqs. (1.4) and (1.9) satisfies the conditions

IV In the Euclidean region ($z = x + iy$, $x > 0$)

$$\tilde{V}(x) > 0; \quad (1.14)$$

$$V \quad D_0(0) = \frac{1}{(2\pi)^4} \int \frac{d^4 k \tilde{V}(\ell^2 k^2)}{m^2 + k^2} < \infty. \quad (1.15)$$

Secondly, we shall not take the Lagrangian in the normal form.

Within the framework of the nonlocal theory this can be done because

$D_0(0) < \infty$, in accordance with Eq. (1.15). The consequence of this assumption will become clear below.

2. Existence conditions for $S_n(x_1, \dots, x_2)$

We shall assume that the function $U(\phi)$ in the interaction Lagrangian

is analytic near $\phi = 0$, so that the series

$$U(a) = \sum_{m=0}^{\infty} \frac{u_m}{m!} a^m \quad (2.1)$$

converges in the circle $|a| < \rho$. The coefficients u_m are real

numbers, and the function $U(a)$ can be written in the form

$$U(a) = \int e^{ia \cdot \zeta} d\sigma(\zeta), \quad (2.2)$$

where $\sigma(\zeta)$ is a certain fully additive measure in the complex plane of ζ ,

such that the integral

$$\int e^{\rho |\zeta|} |d\sigma(\zeta)| < \infty, \quad (2.3)$$

will converge, where ρ is the convergence radius for (2.1).

For the functions $S_n(x_1, \dots, x_n)$ in Eq. (1.10) we then obtain

$$\begin{aligned} S_n(x_1, \dots, x_n) &= \exp \left\{ -\frac{1}{2} \sum_{i,j=1}^n D_0(x_i - x_j) - \frac{\partial^2}{\partial a_i \partial a_j} \right\} \prod_{j=1}^n U(a_j) = \\ &= \int d\sigma(\zeta_1) \dots \int d\sigma(\zeta_n) e^{i \sum_{j=1}^n a_j \zeta_j} \exp \left\{ -\frac{1}{2} \sum_{i,j=1}^n \zeta_i D_0(x_i - x_j) \zeta_j \right\}. \end{aligned} \quad (2.4)$$

We shall now derive the conditions under which this representation is mathematically justified.

We note first that the quadratic form $\sum_{i,j} a_i D_0(x_i - x_j) a_j$ is positive definite for any real $a_1, \dots, a_n$, since

$$\sum_{i,j=1}^n a_i D_0(x_i - x_j) a_j = \frac{1}{(2\pi)^4} \int \frac{d^4 k \tilde{V}(\ell^2 k^2)}{m^2 + k^2} \left| \sum_{j=1}^n a_j e^{ikx_j} \right|^2 > 0. \quad (2.5)$$

On the other hand, we can readily find the upper bound for this form:

$$\begin{aligned} \sum_{i,j=1}^n a_i D_0(x_i - x_j) a_j &\leq \frac{1}{(2\pi)^4} \int \frac{d^4 k \tilde{V}(\ell^2 k^2)}{m^2 + k^2} (a_1 + \dots + a_n)^2 \leq \\ &\leq n D_0(0) (a_1^2 + \dots + a_n^2), \end{aligned} \quad (2.6)$$

and this estimate cannot be improved. Thus,

$$0 \leq \sum_{i,j=1}^n a_i D_0(x_i - x_j) a_j \leq n D_0(0) (a_1^2 + \dots + a_n^2), \quad (2.7)$$

Using this inequality, and substituting $\zeta_j = \xi_j + i \eta_j$ in Eq. (2.4), we obtain the following estimate:

$$\begin{aligned}
 |S_n(x_1, \dots, x_n)| &\leq \int |d\sigma(\zeta_1)| \dots \int |d\sigma(\zeta_n)| e^{\sum_{j=1}^n a_j \eta_j} \times \\
 &\times \exp \left\{ -\frac{1}{2} \sum_{j=1}^n \xi_j D_0(x_1 - x_j) \xi_j + \frac{1}{2} \sum_{j=1}^n \eta_j D_0(x_1 - x_j) \eta_j \right\} \leq \\
 &< \prod_{j=1}^n \int |d\sigma(\zeta_j)| \exp \left\{ a_j \eta_j + \frac{n}{2} D_0(0) \eta_j^2 \right\}
 \end{aligned} \tag{2.8}$$

Hence it follows immediately that the expression given by Eq. (2.4) is mathematically defined if the integral

$$\int |d\sigma(\zeta)| e^{N \eta^2} < \infty \tag{2.9}$$

converges for any $N > 0$.

We must now find the class of functions $U(a)$ which satisfy Eq. (2.3) and (2.9). Substituting $\alpha \rightarrow a + i\beta$, we obtain

$$\begin{aligned}
 |U(a + i\beta)| &= \left| \int d\sigma(\zeta) e^{i(a+i\beta)(\zeta+i\eta)} \right| \leq \\
 &\leq \int |d\sigma(\zeta)| e^{|a||\eta| + |\beta||\zeta|} < \max_{|\eta|, |\zeta|} e^{|a||\eta| - N|\eta|^2} e^{|\beta||\zeta| - \rho|\zeta|} \int |d\sigma(\zeta)| e^{N^2 \eta^2 + \rho|\zeta|} \\
 &\leq \text{Const} e^{\frac{1}{N} a^2} \max_{|\zeta|} e^{(|\beta| - \rho)|\zeta|}
 \end{aligned} \tag{2.10}$$

for any $N > 0$.

It follows from this estimate that the function $U(\alpha + i\beta)$ should be

1) analytic in the band

$$-\infty < \alpha < \infty, \quad -\rho < \beta < \rho; \quad (2.11)$$

2) it should increase within the band more slowly than any Gaussian exponential, i.e.

$$\lim_{\alpha \rightarrow \pm \infty} |U(\alpha + i\beta)| e^{-\epsilon \alpha^2} = 0 \quad (2.12)$$

for $|\beta| < \rho$ and any $\epsilon > 0$.

For other functions $U(a)$ and, in particular, for functions having singularities on the real axis of a , the representation given by Eq. (2.4) is mathematically undefined.

Thus, for the class of functions $U(a)$ satisfying the above conditions, the S-matrix will be finite, unitary, and macrocausal in each order of the perturbation theory.

It readily follows from Eq. (2.8) that

$$|S_n(x_1, \dots, x_n)| < e^{a(n)}, \quad (2.13)$$

where

$$s(n) > A n^2 \quad (2.14)$$

Hence it also follows that the perturbation theory series diverges fairly rapidly. Powerful methods of summation will be required to obtain the complete S-matrix of the theory.

An analogous result was obtained in ref. 4 in the construction of a finite perturbation-theory series with arbitrary interaction Lagrangians.

3. Convergence of the perturbation-theory series

Among the functions $U(a)$ satisfying the above conditions, a special place is occupied by functions which decrease as $a \rightarrow \pm\infty$, i.e.

$$\lim_{a \rightarrow \pm\infty} U(a) = 0. \quad (3.1)$$

This means that for these functions we have a Fourier representation

$$U(a) = \int_{-\infty}^{\infty} d\xi e^{ia\xi} \bar{U}(\xi), \quad (3.2)$$

where

$$\int_{-\infty}^{\infty} d\xi e^{|\alpha||\xi|} |\bar{U}(\xi)| < \infty \quad (3.3)$$

for $|a| < \rho$.

For such functions we can always find the representation (2.4) for

$$S_n(x_1, \dots, x_n).$$

$$S_n(x_1, \dots, x_n) = \int_{-\infty}^{\infty} d\xi_1 \dots \int_{-\infty}^{\infty} d\xi_n U(\xi_1) \dots U(\xi_n) e^{i \sum_{j=1}^n a_j \xi_j x_j} \times \\ \times \exp \left\{ - \frac{1}{2} \sum_{i,j=1}^n \xi_i D_0(x_i - x_j) \xi_j \right\}. \quad (3.4)$$

According to Eq. (2.7), we have

$$|S_n(x_1, \dots, x_n)| < \left| \int_{-\infty}^{\infty} d\xi | \bar{U}(\xi) | \right|^n \quad (3.5)$$

Thus, in this case the expression given by Eq. (3.4) is well defined and, moreover, Eq. (3.5) shows that the perturbation-theory series converges.

Let us introduce into the expansion of the S-matrix in Eq. (1.9) a function $g(x)$ instead of the coupling constant g . This function characterizes the degree to which the interaction is switched on, and is such that

$$\int d^4x \, g(x) < \infty \quad (3.6)$$

The series

$$S = \sum_{n=0}^{\infty} \frac{1}{n!} \int d^4 x_1 \dots \int d^4 x_n g(x_1) \dots g(x_n) S_n(x_1, \dots, x_n) \quad (3.7)$$

is then majored by the series

$$\begin{aligned} S_M &= \sum_{n=0}^{\infty} \frac{1}{n!} \left\{ \int d^4 x g(x) \int_{-\infty}^{\infty} d\xi |U(\xi)| \right\}^n = \\ &= \exp \left\{ \int d^4 x g(x) \int_{-\infty}^{\infty} d\xi |\tilde{U}(\xi)| \right\}. \end{aligned} \quad (3.8)$$

Consequently, Eq. (3.7) converges absolutely. The quantity

$$\int d^4 x g(x) = g V_4 \quad (3.9)$$

diverges since $V_4 \rightarrow \infty$, where V_4 has the meaning of the four-

dimensional volume for $g(x) \rightarrow g$. In the usual quantum field theory these

divergences are connected with vacuum-vacuum transitions, and are present in

the perturbation-theory series in the form of an unimportant phase factor

(see, for example, ref. 8).

4. Normal form of interaction Lagrangian

In deriving representation (3.4) and estimate (3.5), it was assumed that

$$L_I(x) = g \int_{-\infty}^{\infty} d\xi e^{i\xi\phi(x)} \tilde{U}(\xi) \quad (4.1)$$

was not chosen in a normally ordered form. If, on the other hand, the interaction Lagrangian is chosen as usual, in the normally ordered form

$$L_I(x) = g \int_{-\infty}^{\infty} d\xi : e^{i\xi\phi(x)} : \tilde{U}(\xi) \quad (4.2)$$

then, instead of Eq. (3.4), we obtain

$$S_n(x_1, \dots, x_n) = \int_{-\infty}^{\infty} d\xi_1 \dots \int_{-\infty}^{\infty} d\xi_n \prod_{j=1}^n \tilde{U}(\xi_j) e^{-\frac{1}{2} D_0^{(0)} \xi_1^2 + i a_j \xi_j} \times \\ \times \exp \left\{ -\frac{1}{2} \sum_{i,j=1}^n \xi_i D_0(x_i - x_j) \xi_j \right\}. \quad (4.3)$$

The necessary condition for the convergence of the integrals in Eq. (4.3) is

that the integral

$$\int_{-\infty}^{\infty} d\xi : \tilde{U}(\xi) : e^{-\frac{1}{2} D_0^{(0)} \xi^2} < \infty \quad (4.4)$$

should converge. Hence it follows that the function $U(a)$ in the plane

of the complex variable a should be an integral analytic function of

order $p \leq 2$.

On the other hand, if in the interaction Lagrangian (4.1) we transform to the normal form, we obtain

$$: U(\phi) : = \int_{-\infty}^{\infty} d\xi : e^{i\xi\phi(x)} : e^{-\frac{1}{2} D_0^{(0)} \xi^2} \tilde{U}(\xi). \quad (4.5)$$

Therefore, the normally ordered interaction Lagrangian $:U(a):$ is an integral function of order $p \leq 2$ in the plane of a , even if the Lagrangian $U(a)$ has a singularity in this plane.

Moreover, if the interaction Lagrangian is not chosen in the normal form, then after transition to the normal form in the interaction Lagrangian and in each order of perturbation theory for the S-matrix each term in the perturbation-theory series $S_n(x_1, \dots, x_n)$ will depend on the elementary length l in such a way that

$$\lim_{l \rightarrow 0} S_n(x_1, \dots, x_n) = 0 \quad (4.6)$$

Thus, in the procedure which we have been considering, transition to the point interaction is accompanied by disappearance of the real interaction, while in the usual nonlocal theory with $l \rightarrow 0$ there are the usual ultraviolet divergences of the local quantum field theory.

Conclusion

Thus, for a definite class of essentially nonlinear interaction Lagrangians, we have succeeded in constructing within the framework of

nonlocal quantum field theory an S-matrix in the form of a converging perturbation-theory series in the Euclidean formulation of the theory.

It was shown earlier that the S-matrix would be unitary in each order of perturbation theory after continuation into the physical region. The next problem is to investigate the behavior of the amplitudes for physical processes within the framework of the entire converging series representing the complete S-matrix of the theory.

R e f e r e n c e s

1. F. Gursey, *Nuovo Cim.*, 16, 230 (1960); J. Schwinger, *Phys. Letters*, 24B, (1967) 473; S. Coleman, J. Wess, B. Zumino, *Phys. Rev.*, 177, (1969) 2239; C.G. Callen, S. Coleman, J. Wess, B. Zumino, *Phys. Rev.*, 177 (1969) 2247.
2. J. Schwinger, *Phys. Rev.*, 158 (1967) 1391; S. Weinberg, *Phys. Rev. Letters*, 18 (1967) 188.
3. G.V. Efimov, *ZhETF*, 44 (1963) 2107; E.S. Fradkin, *Nucl. Phys.*, 49 (1963) 624; G.V. Efimov, *Nuovo Cim.*, 32 (1964) 1046; G.V. Efimov, *ZhETF* 48 (1965) 598; G.V. Efimov, *Nucl. Phys.*, 74 (1965) 657; M.K. Volkov, *Commun. Math. Phys.*, 7 (1968) 289.
4. G.V. Efimov, JINR, Preprint P2-4472, Dubna, 1969.
5. R. Delbourgo, A. Salam, J. Strathdee, Preprint Trieste, IC-69-17, 1969; IC-69-28, 1969.
6. G.V. Efimov, *Commun. Math. Phys.*, 5 (1967) 42; G.V. Efimov, *Yadernaya Fizika* 4 (1966) 432; G.V. Efimov, Preprint ITF-68-54, Kiev, 1968.
7. G.V. Efimov, *Commun. Math. Phys.*, 7 (1968) 138; G.V. Efimov, Preprint ITF-68-52, Kiev, 1968.
8. N.N. Bogolyubov and D.V. Shirkov, *Vvedenie v teoriyu kvantovannykh polei* [Introduction to the Theory of Quantized Fields] Gostekhizdat, Moscow, 1957.

Received June 18th, 1969.

TRANSLATION SUPPLIED BY

addis TRANSLATIONS International
 Post Office Box 4097
 Woodside, California 94062
 Telephone: (415) 851-1040